

Weihrauch degrees of finding equilibria in sequential games*

Stéphane Le Roux

Département d'informatique
Université libre de Bruxelles, Belgique
Stephane.Le.Roux@ulb.ac.be

Arno Pauly

Computer Laboratory
University of Cambridge, United Kingdom
Arno.Pauly@cl.cam.ac.uk

We consider the degrees of non-computability (Weihrauch degrees) of finding winning strategies (or more generally, Nash equilibria) in infinite sequential games with certain winning sets (or more generally, outcome sets). In particular, we show that as the complexity of the winning sets increases in the difference hierarchy, the complexity of constructing winning strategies increases in the effective Borel hierarchy.

1 Overview

We consider questions of (non)computability related to infinite sequential games played by any countable number of players. The best-known example of such games are Gale-Stewart games [19], which are two-player win/lose games. The existence of winning strategies in (special cases of) Gale-Stewart games is often employed to show that truth-values in certain logics are well-determined. The degrees of noncomputability of variations of (Borel) determinacy [40] can be studied using our techniques, and several are fully classified.

This work falls within the research programme to study the computational content of mathematical theorems in the Weihrauch lattice, which was outlined by BRATTKA and GHERARDI in [7]. In particular, it continues the investigation of the Weihrauch degrees of operations mapping games to their equilibria started in [46]. There, finding pure and mixed Nash equilibria in two-player games with finitely many actions in strategic form were classified.

One motivation for this line of inquiry is the general stance that solution concepts in game theory can only be convincing if the players are capable of (at least jointly) computing them, taken e.g. in [48]. Even if we allow for some degree of hypercomputation, or are, e.g., willing to tacitly replace actually attaining a solution concept by some process (slowly) converging to it, we still have to reject solution concepts with too high a Weihrauch degree.

The results for determinacy of specific pointclasses that we provide are a refinement of results obtained in reverse mathematics by NEMOTO, MEDSALEM and TANAKA [45]; the first is also a uniformization of a result by CENZER and REMMEL [14]. For some represented pointclass Γ , let $\text{Det}_\Gamma : \Gamma \rightrightarrows \{0,1\}^\mathbb{N}$ be the map taking a Γ -subset A of Cantor space to a (suitably encoded) Nash equilibrium in the sequential two-player game with alternating moves where the first player wins if the induced play is in A , and the second player wins otherwise. Let $\mathcal{A}$ be the closed subsets of Cantor space, and $\mathfrak{D} := \{U \setminus U' \mid U, U' \in \mathcal{A}\}$. Some of our results are:

Theorem. $\text{Det}_\mathcal{A} \equiv_W C_{\{0,1\}^\mathbb{N}}$ and $\text{Det}_\mathfrak{D} \equiv_W C_{\{0,1\}^\mathbb{N}} \star \text{lim}$.

*An extended abstract of this work has appeared in the Proceedings of CiE 2015 [38].

We have two remarks. One, by combining the preceding theorem with the main result of [10], we find that $\text{Det}_{\mathfrak{D}}$ is equivalent to the Bolzano-Weierstrass-Theorem. This may be a bit unexpected in particular seeing that $C_{\{0,1\}^{\mathbb{N}}} \star \text{lim}$ is not (yet) known to contain a plethora of mathematical theorems (unlike, e.g., $C_{\{0,1\}^{\mathbb{N}}}$). Two, we already need to use a limit operator in order to move up one level of the difference hierarchy – rather than being able to move up one level in the Borel hierarchy as one may have expected naively. Thus, this observation may complement Harvey Friedman’s famous result [18] that proving Borel determinacy requires repeated use of the axiom of replacement.

Another group of results is based on inspecting the various results extending Borel determinacy to more general classes of games (and solution concepts) in [34, 35, 36, 37]. If we instantiate these generic results with specific determinacy version as above, we can prove for some of them that they are actually optimal w.r.t. Weihrauch reducibility. We shall state two such classifications explicitly.

Consider two-player sequential games with finitely many *outcomes* and antagonistic (inverse of each other) linear preferences over the outcomes. For any upper set of outcomes w.r.t. some player’s preference let the corresponding set of plays be open or closed. Let $\text{NE}_{\mathcal{O} \cup \mathcal{A}}^{\text{ap}}$ be the operation taking such a game (suitably encoded) and producing a Nash equilibrium. Then:

Theorem. $\text{NE}_{\mathcal{O} \cup \mathcal{A}}^{\text{ap}} \equiv_{\text{W}} C_{\{0,1\}^{\mathbb{N}}} \times \text{LPO}^*$

Those games will even have subgame-perfect equilibria, and we let $\text{SPE}_{\mathcal{O} \cup \mathcal{A}}$ be the operation mapping such games to a subgame-perfect equilibrium.

Theorem. $\text{SPE}_{\mathcal{O} \cup \mathcal{A}} \equiv_{\text{W}} \text{lim}$

Various further classifications are obtained, and adhere to the scheme that algebraic combinations of very common Weihrauch degrees appear, which is already exhibited by our examples above.

2 Fundamentals

2.1 Background on represented spaces

We briefly recall some fundamental concepts on represented spaces following [49], to which the reader shall also be referred for a more detailed presentation. The concept behind represented spaces essentially goes back to WEIHRAUCH and KREITZ [31], the name may have first been used by BRATTKA [3]. A *represented space* is a pair $\mathbf{X} = (X, \delta_X)$ of a set X and a partial surjection $\delta_X : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow X$. A function between represented spaces is a function between the underlying sets. For $f : \mathbf{X} \rightarrow \mathbf{Y}$ and $F : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$, we call F a *realizer* of f (notation $F \vdash f$), iff $\delta_Y(F(p)) = f(\delta_X(p))$ for all $p \in \text{dom}(f\delta_X)$, i.e. if the following diagram commutes:

$$\begin{array}{ccc} \mathbb{N}^{\mathbb{N}} & \xrightarrow{F} & \mathbb{N}^{\mathbb{N}} \\ \downarrow \delta_{\mathbf{X}} & & \downarrow \delta_{\mathbf{Y}} \\ \mathbf{X} & \xrightarrow{f} & \mathbf{Y} \end{array}$$

A map between represented spaces is called *computable* (continuous), iff it has a computable (continuous) realizer. Similarly, we call a point $x \in \mathbf{X}$ *computable*, iff there is some computable

$p \in \mathbb{N}^{\mathbb{N}}$ with $\delta_{\mathbf{X}}(p) = x$. A priori, the notion of a continuous map between represented spaces and a continuous map between topological spaces are distinct and should not be confused!

Given two represented spaces $\mathbf{X}, \mathbf{Y}$ we obtain a third represented space $\mathcal{C}(\mathbf{X}, \mathbf{Y})$ of continuous functions from X to Y by letting $0^n 1 p$ be a $[\delta_X \rightarrow \delta_Y]$ -name for f , if the n -th Turing machine equipped with the oracle p computes a realizer for f . As a consequence of the UTM theorem, $\mathcal{C}(-, -)$ is the exponential in the category of continuous maps between represented spaces, and the evaluation map is even computable (as are the other canonic maps, e.g. currying).

This function space constructor, together with two represented spaces, $\mathbb{N} = (\mathbb{N}, \delta_{\mathbb{N}})$ and $\mathbb{S} = (\{\perp, \top\}, \delta_{\mathbb{S}})$, allows us to obtain a model of ESCARDÓ's synthetic topology [16]. The representation are given by $\delta_{\mathbb{N}}(0^n 10^{\mathbb{N}}) = n$, $\delta_{\mathbb{S}}(0^{\mathbb{N}}) = \perp$ and $\delta_{\mathbb{S}}(p) = \top$ for $p \neq 0^{\mathbb{N}}$. It is straightforward to verify that the computability notion for the represented space $\mathbb{N}$ coincides with classical computability over the natural numbers. The Sierpiński space $\mathbb{S}$ in turn allows us to formalize semi-decidability. The computable functions $f : \mathbb{N} \rightarrow \mathbb{S}$ are exactly those where $f^{-1}(\{\top\})$ is recursively enumerable (and thus $f^{-1}(\{\perp\})$ co-recursively enumerable).

In general, for any represented space $\mathbf{X}$ we obtain two spaces of subsets of $\mathbf{X}$; the space of open sets $\mathcal{O}(\mathbf{X})$ by identifying $f \in \mathcal{C}(\mathbf{X}, \mathbb{S})$ with $f^{-1}(\{\top\})$, and the space of closed sets $\mathcal{A}(\mathbf{X})$ by identifying $f \in \mathcal{C}(\mathbf{X}, \mathbb{S})$ with $f^{-1}(\{\perp\})$. The properties of the spaces of open and closed sets, namely computability of the usual operations, follow from a few particular computable functions on Sierpiński space $\mathbb{S}$ and the fundamental function space properties.

We require further classes of sets (often called pointclasses in this context) as represented spaces. A general approach to this is found in synthetic descriptive set theory suggested in [51]. As shown in [21], that approach yields consistent definitions with the ones used in MOSCHOVAKIS' effective descriptive set theory [42]. Here, we shall just directly provide representations that suffice. Some definitions essentially already appeared in [5] and/or [53].

Definition 1. Given a represented pointclass Γ , we represent $\{A^C \mid A \in \Gamma\}$ by reinterpreting a name for A as a name for A^C . Furthermore, we represent $\{\bigcup_{i \in \mathbb{N}} A_i \mid \forall i \in \mathbb{N} A_i \in \Gamma\}$ by identifying suitable sequences in $\Gamma^{\mathbb{N}}$. Given represented pointclasses Γ_1, Γ_2 , we represent $\{A \setminus B \mid A \in \Gamma_1, B \in \Gamma_2\}$ by identifying the suitable pairs in $\Gamma_1 \times \Gamma_2$.

Note that only the first of the three constructions preserves admissibility of the representations.

We will make use of the jump of a represented space. This is based on the limit operator $\lim : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ defined via $\lim(p)(n) = \lim_{i \rightarrow \infty} p(\langle n, i \rangle)$. We then define $(X, \delta_X)' = (X, \delta_X \circ \lim)$. This is iterated along $\mathbf{X}^{(0)} := \mathbf{X}$ and $\mathbf{X}^{(n+1)} := (\mathbf{X}^{(n)})'$. In [10], the jump was extended to multivalued functions via $(f : \mathbf{X} \rightrightarrows \mathbf{Y})^{[n]} := (f : \mathbf{X}^{[n]} \rightrightarrows \mathbf{Y})$.

2.2 Informal background on infinite sequential games

We use the formal definitions of sequential games and related concepts from [36, 37] and [34]. Informally, given a fixed (w.l.o.g.) set C , we let the players sequentially choose elements in C until an infinite sequence in C^{ω} is generated. Whose turn it is depends on the finite history of choices. The outcome (from a set O) of the game depends on the generated sequence in C^{ω} , and each player may compare outcomes via a binary relation over O , called preference. A strategy of a player is an object that fully specifies what the choice of the player would be for each possible finite history that requires this player to play. A combination of one strategy per player is called a strategy profile and it induces one unique infinite sequence in C^{ω} , and thus

one unique outcome. So, preferences may be lifted from outcomes to strategy profiles. A Nash equilibrium is a profile such that no player can unilaterally change strategies and induce a (new) outcome that he or she prefers over the old one. We also consider a refinement of the concept of a Nash equilibrium, namely subgame-perfect Nash equilibria. Intuitively, a strategy profile is subgame-perfect, if it still forms an equilibrium if the game were started at an arbitrary history.

2.3 Background on infinite sequential games

Our presentation of the required background on infinite sequential games is modelled closely on the corresponding section in [37], and on [34].

It is convenient to introduce the abstract notion of a *game* first, before introducing additional structure later on. A game is a tuple $\langle A, (S_a)_{a \in A}, (\prec_a)_{a \in A} \rangle$ consisting of a non-empty set A of *agents* or *players*, for each agent $a \in A$ a non-empty set S_a of *strategies*, and for each agent $a \in A$ a *preference* relation $\prec_a \subseteq (\prod_{a \in A} S_a) \times (\prod_{a \in A} S_a)$. The generic setting suffices to introduce the notion of a Nash equilibrium: A *strategy profile* $\sigma \in (\prod_{a \in A} S_a)$ is called a Nash equilibrium, if for any agent $a \in A$ and any strategy $s_a \in S_a$ we find $\neg(\sigma \prec_a \sigma_{a \rightarrow s_a})$, where $\sigma_{a \rightarrow s_a}$ is defined by $\sigma_{a \rightarrow s_a}(b) = \sigma(b)$ for $b \in A \setminus \{a\}$ and $\sigma_{a \rightarrow s_a}(a) = s_a$. In words, no agent prefers over a Nash equilibrium some other situation that only differs in her choice of strategy.

Definition 2 (Infinite sequential game, cf. [34, Definition 1.1]). An *infinite sequential game* is an object $\langle A, C, d, O, v, (\prec_a)_{a \in A} \rangle$ complying with the following.

1. A is a non-empty set (of agents).
2. C is a non-empty set (of choices).
3. $d : C^* \rightarrow A$ (assigns a decision maker to each stage of the game).
4. O is a non-empty set (of possible outcomes of the game).
5. $v : C^\omega \rightarrow O$ (uses outcomes to value the infinite sequences of choices).
6. Each $\prec_a$ is a binary relation over O (modelling the preference of agent a).

The intuition behind the definition is that agents take turns to make a choice. Whose turn it is depends on the past choices via the function d (which often will be just alternating play). Over time, the agents thus jointly generate some infinite sequence, which is mapped by v to the outcome of the game. Note that using a single set of actions C for each step just simplifies the notation; a generalization to varying action sets is straightforward. In the present paper, typically A and O will be countable, and C even finite.

The infinite sequential games are linked to abstract games as follows: the agents remain the agents and the strategies of agent a are the functions $s_a : d^{-1}(\{a\}) \rightarrow C$. We can then safely regard a strategy profile as a function $\sigma : C^* \rightarrow C$ whose *induced play* is defined below, where for an infinite sequence $p \in C^\omega$ we let p_n be its n -th value, and $p_{\leq n} = p_{< n+1} \in C^*$ be its finite prefix of length n .

Definition 3 (Induced play and outcome, cf. [34, Definition 1.3]). Let $s : C^* \rightarrow C$ be a strategy profile. The *play* $p = p^\lambda(s) \in C^\omega$ *induced by s starting at $\lambda \in C^*$* is defined inductively through its prefixes: $p_n = \lambda_n$ for $n \leq |\lambda|$ and $p_n := s(p_{< n})$ for $n > |\lambda|$. Also, $v \circ p_\lambda(s)$ is the *outcome induced by s starting at λ* . The play (resp. outcome) induced by s is the play (resp. outcome) induced by s starting at ε .

In the usual way to regard an infinite sequential game as a special abstract game, an agent prefers a strategy profile σ to σ' , iff he prefers the outcome induced by σ to the outcome induced by σ' . And indeed we shall call a strategy profile of an infinite sequential game a Nash equilibrium, iff it is a Nash equilibrium with these preferences. In a certain notation overload, we will in particular use the same symbols for the preferences over strategy profiles and the preferences over outcomes.

We also consider a refinement of the concept of a Nash equilibrium, namely subgame-perfect Nash equilibria. Intuitively, a strategy profile is subgame-perfect, if it still forms an equilibrium if the game were started in an arbitrary position.

As a important special case we consider win/lose games. These are games with two players a, b and two outcomes w_a, w_b , where a prefers w_a to w_b and b prefers w_b to w_a . We say that a wins the game, if outcome w_a is reached, and call the set of all plays that induce outcome w_a as the winning set for a (likewise for b and w_b).

2.4 Background on Weihrauch reducibility

A few years ago several authors (GHERARDI and MARCONE [20], P. [47, 46], BRATTKA and GHERARDI [7]) noticed that a reducibility notion based on previous work by WEIHRAUCH [55, 56] would provide a very interesting setting for a metamathematical inquiry into the computational content of mathematical theorems. The fundamental research programme was outlined in [7], and the introduction in [9] may serve as a recent survey.

Definition 4 (Weihrauch reducibility). Let f, g be multi-valued functions on represented spaces. Then f is said to be *Weihrauch reducible* to g , in symbols $f \leq_W g$, if there are computable functions $K, H : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that $K(\text{id}, GH) \vdash f$ for all $G \vdash g$.

The relation $\leq_W$ is reflexive and transitive. We use $\equiv_W$ to denote equivalence regarding $\leq_W$, and by $<_W$ we denote strict reducibility. By $\mathfrak{W}$ we refer to the partially ordered set of equivalence classes. As shown in [47, 8], $\mathfrak{W}$ is a distributive lattice, and also the usual product operation on multivalued function induces an operation $\times$ on $\mathfrak{W}$. The algebraic structure on $\mathfrak{W}$ has been investigated in further detail in [25, 12].

There are two relevant unary operations defined on $\mathfrak{W}$, both happen to be closure operators. The operation $*$ was introduced in [47, 46] by setting $f^0 := \text{id}_{\mathbb{N}^{\mathbb{N}}}$, $f^{n+1} := f \times f^n$ and then $f^*(n, x) := f^n(x)$. It corresponds to making any finite number of parallel uses of f available. Similarly, the *parallelization* operation $\hat{}$ from [8, 7] makes countably many parallel uses available by $\hat{f}(x_0, x_1, x_2, \dots) := (f(x_0), f(x_1), f(x_2), \dots)$.

We will make use of an operation $\star$ defined on $\mathfrak{W}$ that captures aspects of function composition. Following [10, 11], let $f \star g := \max_{\leq_W} \{f_0 \circ g_0 \mid f \equiv_W f_0 \wedge g \equiv_W g_0\}$. We understand that the quantification is running over all suitable functions f_0, g_0 with matching types for the function composition. It is not obvious that this maximum always exists, this is shown in [12] using an explicit construction for $f \star g$. Like function composition, $\star$ is associative but generally not commutative. We use $\star$ to introduce iterated composition via setting $f^{(0)} := \text{id}_{\mathbb{N}^{\mathbb{N}}}$ and $f^{(n+1)} = f^{(n)} \star f$.

All computable multivalued functions with a computable point in their domain are Weihrauch equivalent, this degree is denoted by 1.

An important source for examples of Weihrauch degrees that are relevant in order to classify theorems are the closed choice principles studied in e.g. [7, 6]:

Definition 5. Given a represented space $\mathbf{X}$, the associated closed choice principle $C_{\mathbf{X}}$ is the partial multivalued function $C_{\mathbf{X}} : \subseteq \mathcal{A}(\mathbf{X}) \rightrightarrows \mathbf{X}$ mapping a non-empty closed set to an arbitrary point in it.

For any uncountable compact metric space $\mathbf{X}$ we find that $C_{\mathbf{X}} \equiv_W C_{[0,1]}$. For well-behaved spaces, using closed choice iteratively does not increase its power, in particular $C_{\mathbb{N}} \star C_{\mathbb{N}} \equiv_W C_{\mathbb{N}}$ and $C_{[0,1]} \star C_{[0,1]} \equiv_W C_{[0,1]}$. Likewise, it was shown that $C_{\mathbb{R}^n} \equiv_W C_{\mathbb{R}^n} \star C_{\mathbb{R}^n} \equiv_W C_{\mathbb{N}} \times C_{[0,1]} \equiv_W C_{\mathbb{N}} \star C_{[0,1]} \equiv_W C_{[0,1]} \star C_{\mathbb{N}}$ for any $n > 0$. Closed choice for $[0, 1]$ and $\{0, 1\}^{\mathbb{N}}$ is incomparable. Furthermore, $C_{\{0,1\}^{\mathbb{N}}} \equiv_W \widehat{C_{\{0,1\}^{\mathbb{N}}}} \equiv_W \widehat{C_{\{0,1\}}}$. The degree $C_{[0,1]}$ is closely linked to WKL in reverse mathematics, while $C_{\mathbb{N}}$ is Weihrauch-complete for functions computable with finitely many mindchanges.

Another typical degree is LPO, which has important representatives such as $\neg : \mathbb{S} \rightarrow \mathbb{S}$, the characteristic function of $0^{\mathbb{N}}$, the characteristic function of 0 in $\mathbb{R}$, $\neq : \{0, 1\}^{\mathbb{N}} \times \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}$ and $\text{IsEmpty} : \mathcal{A}(\{0, 1\}^{\mathbb{N}}) \rightarrow \{0, 1\}$.

Furthermore, we require the degree obtained from the limit operator $\lim : \subseteq \mathbb{N}^{\mathbb{N}} \rightarrow \mathbb{N}^{\mathbb{N}}$. This degree was studied by VON STEIN [54], MYLATZ [43] and BRATTKA [4, 5], with the latter noting in [5] that it is closely connected to the Borel hierarchy. HOYRUP, ROJAS and WEIHRAUCH have shown that $\lim$ is equivalent to the Radon-Nikodym derivative in [27]; while [52] by P. and FOUCHÉ exhibited some constructions of Radon measures that are equivalent to $\lim$. The degree $\lim$ also appears in the context of models of hypercomputation as shown by ZIEGLER [58, 57], and captures precisely the additional computational power that certain solutions to general relativity could provide beyond computability [26]. It is related to the examples above via $C_{\mathbb{N}} \times C_{[0,1]} <_W \lim \equiv_W \lim \times \lim <_W \lim \star \lim$, and $\text{LPO} <_W \text{LPO}^* <_W C_{\mathbb{N}} <_W \lim \equiv_W \widehat{\text{LPO}}$. The (strict) hierarchy $(\lim^{(n)})_{n \in \mathbb{N}}$ plays a very similar role in the Weihrauch degrees as the iterated Halting problems fill in the Turing degrees. We always find $f^{[n]} \leq_W f \star \lim^{(n)}$, and for many natural functions f , equivalence holds, since $(f \times \text{id})^{[n]} \equiv_W f \star \lim^{(n)}$.

2.5 Defining the problems of interest

Let Γ be a represented pointclass over $\{0, 1\}^{\mathbb{N}}$. In a straightforward fashion, we can obtain a representation of the infinite sequential games with countably many agents, countably many outcomes, sets of choices $C = \{0, 1\}$ and Γ -measurable valuation function $v : \{0, 1\}^{\mathbb{N}} \rightarrow \mathcal{O}$. The representation encodes the number of agents and outcomes available, for each upper set of outcome the Γ -set of plays resulting in it, the map d as a look-up table, and the relations $\prec_a$ as look-up tables. We always assume that the inverse of any preference relation is well-founded (this guarantees that equilibria exist). Using a canonic isomorphism $\{0, 1\}^* \cong \mathbb{N}$, we will pretend that the space of strategy profiles in such a game is $\{0, 1\}^{\mathbb{N}}$.

We now consider the following multivalued functions:

1. Det_{Γ} takes a two-player win/lose game as input, where the first player has a winning set in Γ . Valid outputs are the Nash equilibria, i.e. the pairs of strategies where one strategy is a winning strategy.
2. Win_{Γ} has the same inputs as Det_{Γ} , and decides which player (if any) has a winning strategy.
3. FindWS_{Γ} is the restriction of Det_{Γ} to games where the first player has a winning strategy.
4. NE_{Γ} takes as input a game with countably many players, finitely many outcomes, and linear preferences, where each upper set of outcomes (w.r.t. each player preference) comes from a Γ -set. The valid outputs are the Nash equilibria.

5. $\text{NE}_\Gamma^{\omega o}$ takes as input a game with countably many players, countably many outcomes, and linear preferences, where each upper set of outcomes (w.r.t. each player preference) comes from a Γ -set. The valid outputs are the Nash equilibria.
6. NE_Γ^{ap} is the restriction of NE_Γ to the two-player games with antagonistic preferences (i.e. $\prec_a = \prec_b^{-1}$).
7. SPE_Γ takes as input a two-player game with finitely many outcomes and antagonistic preferences, where each upper set of outcomes comes from a Γ -set. Valid outputs are the subgame perfect equilibria.

We abbreviate $\bar{\Gamma} := \{U^C \mid U \in \Gamma\}$. Some trivial reducibilities between these problems are: $\text{Win}_\Gamma \equiv_W \text{Win}_{\bar{\Gamma}}$, $\text{Det}_\Gamma \equiv_W \text{Det}_{\bar{\Gamma}}$, $\text{FindWS}_\Gamma \leq_W \text{Det}_\Gamma \leq_W \text{NE}_{\Gamma \cup \bar{\Gamma}}^{ap} \leq_W \text{SPE}_{\Gamma \cup \bar{\Gamma}}$ and $\text{NE}_\Gamma^{ap} \leq_W \text{NE}_\Gamma$.

Throughout the paper we assume that Γ is determined (which implies that all operations are well-defined in the first place), closed under rescaling and finite intersection with clopens, and that $\emptyset, \{0, 1\}^\mathbb{N} \in \Gamma$. All such closure properties (including those appearing as conditions in the results) are assumed to hold in a uniformly computable way, e.g. given a name for a set in Γ and a clopen, we can compute a name for the intersection of the set with the clopen. With rescaling we refer to the operation $(w, A) \mapsto \{wp \mid p \in A\} : \{0, 1\}^* \times \Gamma \rightarrow \Gamma$ and its inverse.

2.6 The difference hierarchy

The pointclasses we shall study in particular are the levels of the Hausdorff difference hierarchy. Intuitively, these are the sets that can be obtained as boolean combinations of open sets; and their level denotes the least complexity of a suitable term. Roughly following [30, Section 22.E], we shall recall the definition of the difference hierarchy. We define a function par from the countable ordinals to $\{0, 1\}$ by $\text{par}(\alpha) = 0$, if there is a limit ordinal β and a number $n \in \mathbb{N}$ such that $\alpha = \beta + 2n$; and $\text{par}(\alpha) = 1$ otherwise. For a fixed ordinal α , we let $\mathfrak{D}_\alpha$ be the collection of sets D definable in terms of a family $(U_\lambda)_{\lambda < \alpha}$ of open sets via:

$$x \in D \Leftrightarrow \text{par}(\inf\{\beta \mid x \in U_\beta\}) \neq \text{par}(\alpha)$$

In the preceding formula, we understand that $\inf \emptyset = \alpha$. In particular, $\mathfrak{D}_0 = \{\emptyset\}$ and $\mathfrak{D}_1 = \mathcal{O}$.

For our constructions, a different characterization is more useful, though: For some pointclass Γ , let $\mathfrak{D}(\Gamma) := \{\bigcup_{i \in I} v_i U_i \mid \forall i, j \in I \ v_i \in \{0, 1\}^* \wedge U_i \in \Gamma \wedge v_i \not\leq v_j\}$.

Lemma 6. $\mathfrak{D}_{\alpha+1} = \mathfrak{D}(\overline{\mathfrak{D}_\alpha})$ and, more generally, $\mathfrak{D}_\alpha = \mathfrak{D}\left(\overline{\bigcup_{\lambda < \alpha} \mathfrak{D}_\lambda}\right)$

Proof. The proof proceeds via induction. The base case is $\mathfrak{D}_1 = \mathcal{O} = \mathfrak{D}(\{\{0, 1\}^\mathbb{N}\})$, and straightforward. Now assume that the claim holds for all $\beta < \alpha$.

If $A \in \mathfrak{D}_{\alpha+1}$, then there is a witnessing family $(U_\beta)_{\beta < \alpha+1}$ of open sets. Let $U_\alpha = \bigcup_{i \in I} v_i \{0, 1\}^\mathbb{N}$ with a prefix free family $(v_i)_{i \in I}$. For $\lambda < \alpha$ and $i \in I$, we define an open set $U_\lambda^i := \{y \in \{0, 1\}^\mathbb{N} \mid v_i y \in U_\lambda\}$. Then let $A^i \in \mathfrak{D}_\alpha$ be the set constructed from the family $(U_\lambda^i)_{\lambda < \alpha}$, and let $U^i := (A^i)^C$. Now $A = \bigcup_{i \in I} v_i U^i$ witnesses that $A \in \mathfrak{D}(\overline{\mathfrak{D}_\alpha})$.

If $A \in \mathfrak{D}(\overline{\mathfrak{D}_\alpha})$, then there are witnesses $(v_i)_{i \in I}$ and $(U^i)_{i \in I}$ with $U^i \in \overline{\mathfrak{D}_\alpha}$. The latter is in turn witnessed by families of open sets $(U_\beta^i)_{\beta < \alpha}$. For $\beta < \alpha$, let $U_\beta = \bigcup_{i \in I} v_i U_\beta^i$, this is an open set again. Additionally, let $U_\alpha = \bigcup_{i \in I} v_i \{0, 1\}^\mathbb{N}$. Now the family $(U_\lambda)_{\lambda < \alpha+1}$ witnesses that $A \in \mathfrak{D}_{\alpha+1}$. $\square$

Observation 7. If A_n is in $\mathfrak{D}_\alpha$ for all $n \in \mathbb{N}$, so is $A := \bigcup_{n \in \mathbb{N}} 0^n 1 A_n$.

Proof. For all n the set A_n can be written $\bigcup_{i \in I_n} v_{ni} A_{ni}$, where $v_{ni} \not\leq v_{nj}$ and $A_{ni} \in \mathfrak{D}_{\lambda_{ni}}$ for some $\lambda_{ni} < \alpha$, so $A = \bigcup_{n \in \mathbb{N}, i \in I_n} 0^n 1 v_{ni} A_{ni}$, where $0^n 1 v_{ni} \not\leq 0^m 1 v_{mj}$. $\square$

Corollary 8. If B_n is in $\overline{\mathfrak{D}_\alpha}$ for all $n \in \mathbb{N}$, so is $B := \{0^\mathbb{N}\} \cup \bigcup_{n \in \mathbb{N}} 0^n 1 B_n$.

A fundamental result on the difference hierarchy is the Hausdorff-Kuratowski theorem stating that $\bigcup_{\alpha < \omega_1} \mathfrak{D}_\alpha = \Delta_2^0$ (where ω_1 is the smallest uncountable ordinal), see e.g. [30, Theorem 22.27].

3 The computational content of some determinacy principles

We begin by classifying the simplest non-computable games, namely games where the first player wants to reach some closed set. This classification essentially is a uniform version of a result by CENZER and REMMEL [14].

Theorem 9. $\text{FindWS}_\mathcal{A} \equiv_W \text{Det}_\mathcal{A} \equiv_W C_{\{0,1\}^\mathbb{N}}$.

Proof. $C_{\{0,1\}^\mathbb{N}} \leq_W \text{FindWS}_\mathcal{A}$ Given a closed subset $A \in \mathcal{A}(\{0,1\}^\mathbb{N})$, we can easily obtain the game where only player 1 moves, and player 1 wins iff the induced play falls in A . If A is non-empty, then player 1 has a winning strategy: Play any infinite sequence in A .

$\text{FindWS}_\mathcal{A} \leq_W \text{Det}_\mathcal{A}$ Trivial.

$\text{Det}_\mathcal{A} \leq_W C_{\{0,1\}^\mathbb{N}}$ Given the open winning set of player 2, we can modify the game tree by ending the game once we know for sure that player 2 will win. Now the set of strategy profiles where either player 1 wins and player 2 cannot win, or player 2 wins and player 1 cannot prolong the game, is a closed set effectively obtainable from the game. Moreover, it is non-empty, and any such strategy profile is a Nash equilibrium. $\square$

Proposition 10. $\text{Win}_\mathcal{A} \equiv_W \text{LPO}$

Proof. This follows by combining the constructions from the preceding theorem with the fact that $\text{IsEmpty} : \mathcal{A}(\{0,1\}^\mathbb{N}) \rightarrow \{0,1\}$ is equivalent to LPO. $\square$

We can use the results for $\mathcal{A}$ as the base case for classifying the strength of determinacy for the difference hierarchy.

Lemma 11 ⁽¹⁾. $\text{Det}_{\mathfrak{D}(\Gamma)} \leq_W C_{\{0,1\}^\mathbb{N}} \star (\widehat{\text{Det}_\Gamma \times \text{Win}_\Gamma})$ and $\text{Win}_{\mathfrak{D}(\Gamma)} \leq_W \text{LPO} \star_s \widehat{\text{Win}_\Gamma}$.²

Proof. Let the winning set of the first player in the original game be $\bigcup_{i \in \mathbb{N}} v_i U_i$. We use $\widehat{\text{Det}_\Gamma \times \text{Win}_\Gamma}$ to find out who wins each of the games with winning sets U_i , and a Nash equilibrium for each such game. Let v_{n_i} be the subsequence of the v_n where the first player wins.

Now consider the game where the first player's winning set is the open set $U' = \bigcup_{i \in \mathbb{N}} v_{n_i} \{0,1\}^\mathbb{N}$. The first player is winning this derived game, iff he is winning the original game. Thus, by Proposition 10 the claim $\text{Win}_{\mathfrak{D}(\Gamma)} \leq_W \text{LPO} \star_s \widehat{\text{Win}_\Gamma}$ follows. Due to Theorem 9, we can use

¹This is a generalization of the proof idea for [45, Theorem 3.7] by NEMOTO, MEDSALEM and TANAKA. [45, Theorem 3.7] states that ACA_0 proves determinacy for $\mathfrak{D}(\Sigma_1^0)$.

²Here, $\star_s$ denotes the analogue of $\star$ for *strong Weihrauch reducibility* (cf [10]). Its precise definition is not important in the following, and the statement remains true if $\star_s$ is replaced by $\star$.

$C_{\{0,1\}^{\mathbb{N}}}$ to find a Nash equilibrium of this game. Then we combine this Nash equilibrium with those of the subgames to get a Nash equilibrium of the entire game, by letting both players play their equilibrium strategy in any of the subgames rooted at a v_n , and above those, letting the first player try to reach some v_{n_i} and the second to avoid them. This is the claim $\text{Det}_{\mathfrak{D}(\Gamma)} \leq_W C_{\{0,1\}^{\mathbb{N}}} \star \widehat{\text{Det}_{\Gamma} \times \text{Win}_{\Gamma}}$. $\square$

Observation 12. $\text{Det}_{\bigcup_{n \in \mathbb{N}} \Gamma_n} \equiv_W \coprod_{n \in \mathbb{N}} \text{Det}_{\Gamma_n}$ and $\text{Win}_{\bigcup_{n \in \mathbb{N}} \Gamma_n} \equiv_W \coprod_{n \in \mathbb{N}} \text{Win}_{\Gamma_n}$

We will relate deciding the winner and finding a winning strategy for games induced by sets from some level of the difference hierarchy to the *lessor limited principle of omniscience* and the *law of excluded middle* for Σ_n^0 -formulae of the corresponding level. These principles were studied in [1, 10, 24] (among others). Let $(\Sigma_n^0 - \text{LLPO}) : \subseteq \{0, 1\}^{\mathbb{N}} \times \{0, 1\}^{\mathbb{N}} \rightrightarrows \{0, 1\}$ be defined via $i \in (\Sigma_n^0 - \text{LLPO}) (p_0, p_1)$ iff $\forall k_1 \exists k_2 \dots \nexists k_n p_i(\langle k_1, \dots, k_n \rangle) = 1$ (where $\nexists = \forall$ if n is odd and $\nexists = \exists$ otherwise). Let $(\Sigma_n^0 - \text{LEM}) : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}$ be defined via $(\Sigma_n^0 - \text{LEM}) (p) = 1$ iff $\forall k_1 \exists k_2 \dots \nexists k_n p(\langle k_1, \dots, k_n \rangle) = 1$ and $(\Sigma_n^0 - \text{LEM}) (p) = 0$ otherwise. Then:

Proposition 13. $(\Sigma_{n+1}^0 - \text{LLPO}) \equiv_W \text{LLPO}^{[n]}$ and $(\Sigma_{n+1}^0 - \text{LEM}) \equiv_W \text{LPO}^{[n]}$

Lemma 14. $(\widehat{\Sigma_n^0 - \text{LLPO}}) \leq_W \text{Det}_{\mathfrak{D}_n}$ and $(\Sigma_n^0 - \text{LEM}) \leq_W \text{Win}_{\mathfrak{D}_n}$.

Proof. We will first describe the construction for $(\Sigma_n^0 - \text{LEM}) \leq_W \text{Win}_{\mathfrak{D}_n}$, which will then be reused for the remaining claim. The game structure will only depend on the parameter n , but not on the actual input for $(\Sigma_n^0 - \text{LEM})$. This input acts only on the winning set.

The game works as follows: The second player may pick some $k_1 \in \mathbb{N}$, or refuse to play. If the second player picks a number, then the first player may pick $k_2 \in \mathbb{N}$ or refuse to play. This alternating choice continues until k_{n-1} has been chosen, or a player refuses to pick. A player refusing to pick a number loses. If all numbers are picked, the winner depends on the input p to $\Sigma_n^0 - \text{LEM}$ as follows: If n is even and $\exists k_n p(\langle k_1, \dots, k_n \rangle) = 1$, then player 1 wins. If n is odd, and $\exists k_n p(\langle k_1, \dots, k_n \rangle) = 0$, then player 2 wins. Note that this always describes an open component U_{picked} of the winning set of the respective player.

Furthermore, note that the set of plays U_j where a value for k_j was chosen is always an open set. Now the condition that the second player refused to pick first is $U_1^C \cup (U_2 \cap U_3) \cup (U_4 \cap U_5) \cup \dots$. This makes for a winning set in $\mathfrak{D}_n$, as required. If player 1 has a winning strategy in the game, the answer to $(\Sigma_n^0 - \text{LEM}) (p)$ is 1, if player 2 wins, it is 0.

The game for the reduction $(\widehat{\Sigma_n^0 - \text{LLPO}}) \leq_W \text{Det}_{\mathfrak{D}_n}$ adds two layers above the game discussed before. First, player 2 picks an index $j \in \mathbb{N}$ of one of the input pairs $\langle p_1^0, p_1^1 \rangle, \langle p_2^0, p_2^1 \rangle, \dots$ of $(\Sigma_n^0 - \text{LLPO})$, and loses the game if he refuses to pick. Then player 1 picks $i \in \{0, 1\}$, and they play the game above on p_j^i . The extra layers do not impact the complexity of the winning set, in particular since the first two natural numbers are both to be chosen by player 2. The map $j \mapsto i$ extractable from player 1's winning strategy is a valid solution to $(\Sigma_n^0 - \text{LLPO})$. $\square$

Theorem 15. $\text{Det}_{\mathfrak{D}_{n+1}} \equiv_W C_{\{0,1\}^{\mathbb{N}}}^{[n]}$ and $\text{Win}_{\mathfrak{D}_{n+1}} \equiv_W \text{LPO}^{[n]}$.

Proof. Note that $\widehat{\text{LLPO}^{[n]}} \equiv_W \widehat{\text{LLPO}}^{[n]}$ and $\widehat{\text{LPO}^{[n]}} \equiv_W \lim^{(n+1)}$. One direction of the equivalences is provided by Lemma 14 (while taking into consideration Proposition 13). The other direction is shown by induction. The base case is provided by Theorem 9 and Proposition 10. The induction step uses Lemma 11. $\square$

The main obstacle for extending Theorem 15 to the transfinite levels of the difference hierarchy lies in the absence of a standard representation for the space of countable ordinals, which would be required to formulate the statement in the first place. This problem is (partially) addressed in [50]. In the mean time, we can give part of the result for the ω -th level:

Corollary 16. $\text{Det}_{\mathfrak{D}_\omega} \leq_W C_{\{0,1\}^\mathbb{N}} \star \left(\widehat{\prod_{n \in \mathbb{N}} \lim^{(n)}} \right)$ and $\text{Win}_{\mathfrak{D}_\omega} \leq_W \text{LPO} \star \left(\widehat{\prod_{n \in \mathbb{N}} \lim^{(n)}} \right)$

Knowing the Weihrauch degree of a mapping entails some information about the Turing degrees of outputs relative to the Turing degrees of inputs, this was explored in e.g. [8, 10, 9, 48]. Thus, we can obtain the following corollaries:

Corollary 17. Any computable game with a winning condition in $\mathfrak{D}_{n+1}$ has a winning strategy s such that s' is computable relative to $\emptyset^{(n+1)}$, and there is a computable game of this type such that any winning strategy computes a PA-degree relative to $\emptyset^{(n)}$.

Corollary 18. Let $(G_i)_{i \in \mathbb{N}}$ be an effective enumeration of computable games with winning conditions in $\mathfrak{D}_{n+1}$, and define $w \in \{0,1\}^\mathbb{N}$ via $w(i) = 1$ iff the first player has a winning strategy in G_i . Then $w \leq_T \emptyset^{(n+1)}$, and there is an enumeration $(G_i)_{i \in \mathbb{N}}$ such that $w \equiv_T \emptyset^{(n+1)}$.

Corollary 19. There is a Σ_{n+1}^0 -measurable function mapping games with winning conditions in $\mathfrak{D}_n$ to winning strategies, but no Σ_n^0 -measurable such function.

Proof. As shown in [5], the map $\lim^{(n)}$ is Weihrauch-complete for the (effectively) Σ_{n+1}^0 -measurable functions. From Theorem 15 we may conclude that $\lim^{(n)} <_W \text{Det}_{\mathfrak{D}_{n+1}} <_W \lim^{(n+1)}$. $\square$

Before ending this section, we shall make explicit a feature of the constructions used above:

Proposition 20. Consider a pointclass Γ that is closed under the operation $(A_n)_{n \in \mathbb{N}} \mapsto (\{01^\mathbb{N}\} \cup \bigcup_{n \in \mathbb{N}} 1^n 0 A_n)$. Then:

$$\text{FindWS}_\Gamma \equiv_W \widehat{\text{FindWS}_\Gamma}$$

Proof. Given a sequence of games, we construct a single combined game as follows: The second player picks a natural number, and loses the game if he fails to do so. If a number is chosen, the players proceed to the game with the corresponding index. If the first player has winning strategies in all input games, he can win the combined game by playing their combination – and only by doing so. $\square$

4 The complexity of equilibrium transfer

In [34, 35, 36, 37], various results were provided that transfer Borel determinacy (or, somewhat more general, determinacy for some pointclass), to prove the existence of Nash equilibria (and sometimes even subgame-perfect equilibria) in multi-player multi-outcome infinite sequential games. In this section, we shall inspect those constructions and extract Weihrauch reductions from them.

In [35], the first author gave a very general construction that allows to extend determinacy of win/lose games to the existence of Nash equilibria for two-player games of the same type. For brevity, we only consider the strength of the toy example from [35] here:

Theorem 21 (Equilibrium transfer). $\text{NE}_\Gamma^{ap} \leq_W \text{Det}_\Gamma^* \times \text{Win}_\Gamma^*$.

Proof. For any upper set of outcomes (for either players preferences), we construct the win/lose derived game where that player wins, iff he enforces the set, and loses otherwise. There are finitely many such games, so we can use Win_Γ^* to decide which are won and which are lost. As shown in [35], there will be a combination of upper sets of outcomes for each player, such that if both players enforce their upper set, this forms a Nash equilibrium. We use Det_Γ^* to compute Nash equilibria for all derived games in parallel, and then simply select the suitable strategies. $\square$

Techniques suitable for multiplayer sequential games were then introduced in [34], again by the first author. The computational content differs somewhat depending on whether there are finitely many different outcomes, or countably many.

Theorem 22 (Constructing Nash equilibria). $\text{NE}_\Gamma \leq_W \widehat{\text{Win}_\Gamma} \times \widehat{\text{Det}_\Gamma}$,
 $\text{NE}_\Gamma^{\omega_o} \leq_W (\lim \star \widehat{\text{Win}_\Gamma}) \times \widehat{\text{Det}_\Gamma}$

Proof. Let us first prove the first statement. Let us consider a game with finitely many players. (Considering countably many players would be possible, but it would reduce to the finite case since there are only finitely many possible preference relations.) For each node of the game and each upper interval of the preference of the player owning the node (that is, for countably many cases) let us do two things in parallel: on the one hand, invoke Win_Γ and ask who is winning the win/lose subgame rooted at the node where all the opponents of the node owner team up and try to yield an outcome outside of the given preference-upper interval; on the other hand, invoke Det_Γ and obtain a Nash equilibrium for that same game. We claim that this information suffices to computably build a Nash equilibrium for the original game along the proof from [34]: indeed, the *best guarantee* [34, Definition 2.5] of a player, which is the smallest preference-upper interval that the player can enforce, can be computed since we already know who wins the relevant derived games thanks to Win_Γ ; the existential witness from [34, Lemma 2.6] is a Nash equilibrium that has been already computed by Det_Γ ; which is enough to deepen the guarantee as in [34, Definition 2.7] and build the strategy profile σ [34, Lemma 2.8]; and finally, the threats that [34, Theorem 2.9] attaches along the play of σ are given by the Nash equilibria that have been already computed by Det_Γ .

Let us now consider the similar, second statement. At each node, the associate games are now countably many, so knowing who wins each of them no longer suffices to compute the best guarantee of the node owner. To this purpose we use one $\lim$ operator per node, and the rest follows as in the finite case above, so $\text{NE}_\Gamma^{\omega_o} \leq_W (\lim \star \widehat{\text{Win}_\Gamma}) \times \widehat{\text{Det}_\Gamma}$, and the claim follows since $\lim \equiv_W \widehat{\lim}$. $\square$

A further improvement on the techniques in [34] were provided by the authors in [36, 37]. These techniques in particular suffice to prove the existence of subgame-perfect equilibria in antagonistic games (this implies two players and finitely many outcomes).

Theorem 23. $\text{SPE}_\Gamma \leq_W \widehat{\text{Win}_\Gamma} \times \widehat{\text{Det}_\Gamma}$

Proof. Like in the proof of Theorem 22 let us ask at every node who wins the derived games and ask for a corresponding Nash equilibrium. These two pieces of information together provide the existential witness needed in the third condition of [37, Lemma 16], so by the recursive construction in the proof of this lemma, its conclusion follows computably. Let us invoke this conclusion once for each of the two players and combine the obtained strategies into a strategy

profile, which is a subgame-perfect equilibrium by determinacy assumptions. (See [37, Lemma 17] for the details.) $\square$

5 Deciding the winner and finding Nash equilibria


The results in Section 3 show that for many concrete examples of Γ , the problem Det_Γ is inherently multivalued, i.e. not equivalent to any functions between admissible spaces. On the other hand, the upper bounds provided in Section 4 all include Win_Γ , which is of course single-valued. In the current section, we will explore some converse reductions, from deciding the winner to finding Nash equilibria. This generally requires some (rather tame) requirements on the pointclasses involved.

Lemma 24. Let Γ be obtained by Γ_1 by first closing under finite union, rescaling and union with clopens; and then adding complements. Then:

$$\text{Win}_{\Gamma_1}^* \leq_w \text{NE}_\Gamma^{ap}$$

Proof. We first informally describe the construction. Given n win/lose games, the first player starts by announcing which of these games she believes she can win. Then the second player can choose one of the listed games to play. If the first player did not claim any winnable games, the game ends and the outcome is 0. If the first player claimed to be able to win k out of n games, then the outcomes of the games subsequently chosen by the second player are scaled up to $k, -k$. Thus, the first player has every reason to list precisely those games she can actually win: If she would not list a game she could win, she trades payoff $k - 1$ for payoff k . If she lists a game she cannot win, the second player will subsequently chose and win it, and then the first player is punished by $-k$.

The following depicts the construction in case of two input games:




It remains to argue that the resulting game actually is a valid input to NE_Γ^{ap} . We need to show that any upper set of outcomes is associated with a set of plays belonging to Γ . For $k > 0$, the upper set is a finite union of rescaled sets from Γ_1 , which by assumption is a member of Γ . For $k = 0$, to the former we add the clopen set of plays resulting in 0. For $k < 0$, we add additional clopens for those subgames played with stakes less than k . $\square$

Lemma 25. Let Γ be closed under taking unions with Γ_1 and $\overline{\Gamma_1}$. Then:

$$\text{NE}_\Gamma^{ap} \times \text{FindWS}_{\Gamma_1} \equiv_w \text{NE}_\Gamma^{ap}$$

Proof. As clearly $1 \leq_W \text{FindWS}_\Gamma$, we only need to show $\text{NE}_\Gamma^{ap} \times \text{FindWS}_\Gamma \leq_W \text{NE}_\Gamma^{ap}$. Let G_0 be the input game to NE_Γ^{ap} on the left, and G_1 be the input game to FindWS_Γ . If Ω is the (twice ordered) set of outcomes used in G_0 , then we use the outcome set $\Omega \cup \{0, 1\}$ for the newly constructed game, and extend the preferences by $0 \prec_a o \prec_a 1$ and $1 \prec_b o \prec_b 0$ for each $o \in \Omega$. The game tree looks as follows: The second player can choose whether to play in G_1 for outcomes 0 (second player wins) and 1 (second player loses), or to play in G_0 for the original outcomes:



In any Nash equilibrium of this game the second player is choosing to play in G_0 , both players play a Nash equilibrium inside G_0 , and the first player is using a winning strategy inside G_1 . Thus, all desired information can be recovered.

In order for the constructed game to be valid for NE_Γ^{ap} , we need that any upper set from Γ is closed under union with a winning set of player 1. $\square$

Corollary 26. Let Γ_1 and Γ simultaneously satisfy the criteria of the two preceding lemmata. Then:

$$\text{FindWS}_{\Gamma_1}^* \times \text{Win}_{\Gamma_1}^* \leq_W \text{NE}_\Gamma^{ap}$$

Unfortunately, the restrictions on Γ , Γ_1 in place in Lemma 25 (and subsequently Corollary 26) are too strong for the application we have in mind. The result of Corollary 26 can be obtained with weaker conditions though:

Lemma 27. Let Γ be obtained by Γ_1 by first closing under finite union, rescaling and union with clopens; and then adding complements. Then:

$$\text{FindWS}_{\Gamma_1}^* \times \text{Win}_{\Gamma_1}^* \leq_W \text{NE}_\Gamma^{ap}$$


Proof. The reduction directly combines the constructions in Lemma 24 and Lemma 25. We only need to argue that the weaker condition on Γ_1 and Γ suffices to have the valuation Γ -measurable in the resulting game. For this, note that the same reasoning as in Lemma 24 applies, with the addition of a Γ_1 set above the other outcomes, and a $\bar{\Gamma}_1$ set below. $\square$

If we have access to subgame perfect equilibria (and are in a context where they are guaranteed to exist), then we can even decide the winner of countably many games in parallel:

Lemma 28. Let Γ_1 contain the closed sets and be closed under finite unions and the operation $(A_n)_{n \in \mathbb{N}} \mapsto (\{0^\mathbb{N}\} \cup \bigcup_{n \in \mathbb{N}} 0^n 1 A_n)$. Let Γ be obtained from Γ_1 by closing under complements. Then:

$$\widehat{\text{Win}_{\Gamma_1}} \leq_W \text{SPE}_\Gamma$$

Proof. The input to $\widehat{\text{Win}_{\Gamma_1}}$ is a sequence of games $(G_i)_{i \in \mathbb{N}}$ with payoffs 0 and 1. From these, we shall construct a single game G with payoffs 0, 1 and $\frac{1}{2}$. In G , the first player can move right as often as he desires. If he moves left once after having gone right n times, then he is faced with the choice of moving left again and then playing G_n against the second player, or to move right to receive a payoff of $\frac{1}{2}$. If player 1 always moves right, he receives a payoff of 1, as in the picture below.



In a subgame perfect equilibrium, the first player has to choose optimally when deciding between playing G_n and receiving a guaranteed payoff of $\frac{1}{2}$ – which means playing G_n iff he can win it. Thus, any subgame perfect equilibrium of G allows us to decide the winner of each G_n .

It only remains to show that the outcome sets of G fall into Γ . For each $i \in \mathbb{N}$, let $U_i \in \Gamma_1$ be the winning set of the first player in G_i . Then the preimage of $\{1\}$ is $\{1\}^{\mathbb{N}} \cup \bigcup_{i \in \mathbb{N}} 1^i 00 U_i$, thus in Γ_1 . The preimage of $\{\frac{1}{2}\}$ is an open set, and thus in Γ together with the preimage of $\{1, \frac{1}{2}\}$. As Γ is closed under complements, it then contains all relevant preimages. $\square$

6 General games with concrete pointclasses

The general constructions put together with the classifications for specific pointclasses allow us to obtain some concrete Weihrauch degrees. First, we shall see that moving from a win/lose game with closed and open outcomes to a two-player game with several outcomes just complicates the operation of finding Nash equilibria by finitely many uses of LPO in parallel:

Theorem 29. $\text{NE}_{\mathcal{O} \cup \mathcal{A}}^{\text{ap}} \equiv_{\text{W}} \text{C}_{\{0,1\}^{\mathbb{N}}} \times \text{LPO}^*$

Proof. For the reduction $\text{NE}_{\mathcal{O} \cup \mathcal{A}}^{\text{ap}} \leq_{\text{W}} \text{C}_{\{0,1\}^{\mathbb{N}}} \times \text{LPO}^*$, instantiate Theorem 21 with the results from Theorem 9 and Proposition 10.

For the other direction, note that $\text{FindWS}_{\mathcal{A}} \equiv_{\text{W}} \text{FindWS}_{\mathcal{O} \cup \mathcal{A}} \equiv_{\text{W}} \text{C}_{\{0,1\}^{\mathbb{N}}}$ as in Theorem 9; and that $\Gamma_1 := \mathcal{A}$ and $\Gamma := \mathcal{O} \cup \mathcal{A}$ satisfy the requirements of Lemma 27, which then provides the desired result. $\square$

The result can actually be strengthened into the following (by noting that the second game constructed in Lemma 14 is always won by the first player):

Theorem 30. $\text{NE}_{\mathfrak{D}_{n+1} \cup \overline{\mathfrak{D}_{n+1}}}^{\text{ap}} \equiv_{\text{W}} \text{C}_{\{0,1\}^{\mathbb{N}}}^{[n]} \times (\text{LPO}^{[n]})^*$

If one wishes to have subgame-perfect equilibria instead of mere Nash equilibria, then countably many uses of LPO become necessary, and the problem becomes equivalent to $\lim$. Note that as long as there are at least three distinct outcomes, the number of outcomes has no further impact on the Weihrauch degree (due to the nature of the construction used to prove Lemma 28) – unlike the situation in Theorem 29, where the number of outcomes is related to the number of times that LPO is used.

Theorem 31. $\text{SPE}_{\mathfrak{D}_n \cup \overline{\mathfrak{D}_n}} \equiv_{\text{W}} \lim^{(n)}$

Proof. For $\text{SPE}_{\mathfrak{D}_n \cup \overline{\mathfrak{D}_n}} \leq_{\text{W}} \lim^{(n)}$, instantiate Theorem 23 with the results from Theorem 15, and note that $\text{LPO}^{(n)} \equiv_{\text{W}} \lim^{(n)}$ and $\text{C}_{\{0,1\}^{\mathbb{N}}}^{[n]} \leq_{\text{W}} \lim^{(n+1)}$.

For the other direction, we use Lemma 28 (applicable by Corollary 8) together with Proposition 10. $\square$

Regarding Theorem 22, we do not (yet?) have matching lower bounds for any particular pointclass. The gap is exemplified by the following:

Corollary 32. $C_{\{0,1\}^{\mathbb{N}}} \times \text{LPO}^* \leq_W \text{NE}_{\mathcal{O} \cup \mathcal{A}} \leq_W \text{lim}$

Proof. The first reduction follows by Theorem 29. The second reduction follows by instantiating Theorem 22 with the results from Theorem 9 and Proposition 10. $\square$

7 Conclusions and Outlook

With Theorem 15, we have shown that the computational strength of determinacy provides a tight connection between the difference hierarchy and the Borel hierarchy (in form of Corollary 19). Note that winning sets from the difference hierarchy correspond to Boolean combinations of reachability and safety conditions. Corollary 18 then provides an upper bound and a worst case for corresponding decidability questions for logic. Theorem 15 also shows that the computational powers of the players required to find a winning strategy vastly exceeds the computational power required to determine the outcome, thus casting doubt on the adequateness of winning strategies (or Nash equilibria) as adequate solution concepts for infinite sequential games³.

The results in Section 4 contrasted with those in Section 6 essentially show that the proofs in [34, 35, 36, 37] are not too wasteful from a constructive perspective – i.e. the constructions employed are not far less constructive than the theorems proven with them.

There are several immediate avenues for extending the work presented here: The restriction to finite action sets (i.e. finitely branching trees) can mostly be lifted without a significant impact on the proof techniques. Note though that the concrete Weihrauch degrees would change drastically, as in Theorem 9 we would need to replace $C_{\{0,1\}^{\mathbb{N}}}$ by $C_{\mathbb{N}^{\mathbb{N}}}$, with the latter residing in a less explored part of the Weihrauch lattice. The results in [36, 37] are more general than covered here, too (with the same proof complexity). A notion of a product of sequential games could, similarly to the use of a products of bimatrix games in [46] (see also [29, 28]), be used to obtain some results on the products on the corresponding Weihrauch degrees. As in [36, 37], one could extend Theorem 15 to games with real-valued payoff functions of prescribed *level* (introduced by HERTLING in [22, 23], see also [47, 13]).

The study of the strength of determinacy for particular pointclasses in reverse mathematics presumably offers further proofs adaptable into the framework of Weihrauch reducibility, e.g. [44, 41, 15].

Another avenue to explore is the connection to the *basic games* introduced by LACHLAN [33] to formalize some meta-observations on the study of recursively enumerable sets. KUMMER showed that the basic games are equivalent to the computable Gale Stewart games in the difference hierarchy over Σ_2^0 [32]. A natural restriction of the basic games can easily be shown to produce computable Gale-Stewart games in the difference hierarchy over Σ_1^0 as studied in this paper. This raises the question whether here another equivalence can be obtained, which would then allow us the transfer of our classifications to the setting of basic games⁴.

³Then of course, infinite sequential games could justifiably be deemed unrealistic anyway.

⁴This question was originally raised by an anonymous referee.

A precise understanding of the Weihrauch degree of determinacy for various pointclasses is related to determinacy questions of games with transfinite length (as studied in e.g. [39, 17]). For example, Corollary 19 together with Borel determinacy implies that games of length $\omega + \omega$ and winning sets from the finite levels of the difference hierarchy are determined.

Further afield, understanding the Weihrauch degrees of determinacy principles may be a contribution to the development of descriptive set theory in computational/category-theoretical terms as suggested in [51]. In order to deal with determinacy beyond the Borel sets, one may have to adopt the *extended Weihrauch-degrees* recently suggested by BAUER and YOSHIMURA [2] as framework.

References

- [1] Yohji Akama, Stefano Berardi, Susumu Hayashi & Ulrich Kohlenbach (2004): *An arithmetical hierarchy of the law of excluded middle and related principles*. In: *19th IEEE Symposium on Logic in Computer Science (LICS 2004)*, pp. 192–201.
- [2] Andrej Bauer & Kazuto Yoshimura (2014): *The Weihrauch lattice is too small*. CCA 2014.
- [3] Vasco Brattka (1996): *Recursive characterization of computable real-valued functions and relations*. *Theoretical Computer Science* 162, pp. 45–77.
- [4] Vasco Brattka (1999): *Computable Invariance*. *Theoretical Computer Science* 210, pp. 3–20.
- [5] Vasco Brattka (2005): *Effective Borel measurability and reducibility of functions*. *Mathematical Logic Quarterly* 51(1), pp. 19–44, doi:10.1002/malq.200310125.
- [6] Vasco Brattka, Matthew de Brecht & Arno Pauly (2012): *Closed Choice and a Uniform Low Basis Theorem*. *Annals of Pure and Applied Logic* 163(8), pp. 968–1008, doi:10.1016/j.apal.2011.12.020.
- [7] Vasco Brattka & Guido Gherardi (2011): *Effective Choice and Boundedness Principles in Computable Analysis*. *Bulletin of Symbolic Logic* 1, pp. 73 – 117, doi:10.2178/bsl/1294186663. ArXiv:0905.4685.
- [8] Vasco Brattka & Guido Gherardi (2011): *Weihrauch Degrees, Omniscience Principles and Weak Computability*. *Journal of Symbolic Logic* 76, pp. 143 – 176. ArXiv:0905.4679.
- [9] Vasco Brattka, Guido Gherardi & Rupert Hölzl (2015): *Probabilistic computability and choice*. *Information and Computation* 242, pp. 249 – 286, doi:http://dx.doi.org/10.1016/j.ic.2015.03.005. Available at <http://www.sciencedirect.com/science/article/pii/S0890540115000206>. ArXiv 1312.7305.
- [10] Vasco Brattka, Guido Gherardi & Alberto Marcone (2012): *The Bolzano-Weierstrass Theorem is the Jump of Weak König’s Lemma*. *Annals of Pure and Applied Logic* 163(6), pp. 623–625, doi:10.1016/j.apal.2011.10.006. Also arXiv:1101.0792.
- [11] Vasco Brattka, Stéphane Le Roux & Arno Pauly (2012): *On the Computational Content of the Brouwer Fixed Point Theorem*. In S.Barry Cooper, Anuj Dawar & Benedikt Löwe, editors: *How the World Computes, Lecture Notes in Computer Science* 7318, Springer Berlin Heidelberg, pp. 56–67, doi:10.1007/978-3-642-30870-3_7.
- [12] Vasco Brattka & Arno Pauly: *On the algebraic structure of Weihrauch degrees*. forthcoming.
- [13] Matthew de Brecht (2014): *Levels of discontinuity, limit-computability, and jump operators*. In Vasco Brattka, Hannes Diener & Dieter Spreen, editors: *Logic, Computation, Hierarchies*, de Gruyter, pp. 79–108. ArXiv 1312.0697.
- [14] Douglas Cenzer & Jeffrey Remmel (1992): *Recursively presented games and strategies*. *Mathematical Social Sciences* 24(2–3), pp. 117 – 139, doi:http://dx.doi.org/10.1016/0165-4896(92)90059-E.

- [15] Naohi Eguchi (2014): *Infinite Games in the Cantor Space over Admissible Set Theories*. In Kojiro Higuchi, editor: *Proc. of Computability Theory and Foundations of Mathematics*.
- [16] Martin Escardó (2004): *Synthetic topology of datatypes and classical spaces*. *Electronic Notes in Theoretical Computer Science* 87.
- [17] Deborah Sue Fraker (2001): *The Dichotomy in the Determinacy of certain two-person infinite games with moves from $\{0, 1\}$* . Master's thesis, University of Nevada, Las Vegas.
- [18] Harvey Friedman (1971): *Higher set theory and mathematical practice*. *Annals of Mathematical Logic* 2(3), pp. 325–357, doi:10.1016/0003-4843(71)90018-0.
- [19] D. Gale & F.M. Stewart (1953): *Infinite games with perfect information*. In: *Contributions to the theory of games*, *Annals of Mathematical Studies* 28, Princeton University Press, pp. 245–266.
- [20] Guido Gherardi & Alberto Marcone (2009): *How uncomputable is the separable Hahn-Banach theorem?* *Notre Dame Journal of Formal Logic* 50(4), pp. 393–425, doi:10.1215/00294527-2009-018.
- [21] Vassilios Gregoriades, Tamás Kispéter & Arno Pauly (2014): *A comparison of concepts from computable analysis and effective descriptive set theory*. arXiv:1401.3325.
- [22] Peter Hertling (1996): *Unstetigkeitsgrade von Funktionen in der effektiven Analysis*. Ph.D. thesis, Fernuniversität, Gesamthochschule in Hagen.
- [23] Peter Hertling & Klaus Weihrauch (1994): *Levels of degeneracy and exact lower complexity bounds*. In: *6th Canadian Conference on Computational Geometry*, pp. 237–242.
- [24] Kojiro Higuchi & Takayuki Kihara (2014): *Inside the Muchnik degrees I: Discontinuity, learnability and constructivism*. *Annals of Pure and Applied Logic* 165(5), pp. 1058 – 1114, doi:http://dx.doi.org/10.1016/j.apal.2014.01.003.
- [25] Kojiro Higuchi & Arno Pauly (2013): *The degree-structure of Weihrauch-reducibility*. *Logical Methods in Computer Science* 9(2), doi:10.2168/LMCS-9(2:2)2013.
- [26] Mark L. Hogarth (1992): *Does General Relativity Allow an Observer to View an Eternity in a Finite Time?* *Foundations of Physics Letters* 5, pp. 173–181.
- [27] Mathieu Hoyrup, Cristóbal Rojas & Klaus Weihrauch (2012): *Computability of the Radon-Nikodym Derivative*. *Computability* 1(1), pp. 3–13, doi:10.3233/COM-2012-005.
- [28] Xiang Jiang & Arno Pauly (2012): *Efficient Decomposition of Bimatrix Games*. <http://arxiv.org/abs/1212.6355>.
- [29] Xiang Jiang & Arno Pauly (2014): *Efficient Decomposition of Bimatrix Games (Extended Abstract)*. In Fabio Mogavero, Aniello Murano & Moshe Y. Vardi, editors: *Proceedings of Strategic Reasoning 2014, EPTCS* 146.
- [30] A.S. Kechris (1995): *Classical Descriptive Set Theory*. *Graduate Texts in Mathematics* 156, Springer.
- [31] Christoph Kreitz & Klaus Weihrauch (1985): *Theory of Representations*. *Theoretical Computer Science* 38, pp. 35–53.
- [32] Martin Kummer (2006): *The complexity of recursion theoretic games*. *Transactions of the AMS* 358, pp. 59–86, doi:http://dx.doi.org/10.1090/S0002-9947-05-04074-2.
- [33] A. H. Lachlan (1970): *On Some Games Which Are Relevant to the Theory of Recursively Enumerable Sets*. *Annals of Mathematics* 91(2), pp. 291–310.
- [34] Stéphane Le Roux (2013): *Infinite Sequential Nash Equilibria*. *Logical Methods in Computer Science* 9(2).
- [35] Stéphane Le Roux (2014): *From winning strategy to Nash equilibrium*. *Mathematical Logic Quarterly* 60(4-5), pp. 354–371, doi:10.1002/malq.201300034. Available at <http://dx.doi.org/10.1002/malq.201300034>. ArXiv 1203.1866.

- [36] Stéphane Le Roux & Arno Pauly (2014): *Infinite Sequential Games with Real-valued Payoffs*. In: *CSL-LICS '14*, ACM, pp. 62:1–62:10, doi:10.1145/2603088.2603120. Available at <http://doi.acm.org/10.1145/2603088.2603120>.
- [37] Stéphane Le Roux & Arno Pauly (2014): *Infinite sequential games with real-valued payoffs*. arXiv:1401.3325. Available at <http://arxiv.org/abs/1401.3325>.
- [38] Stéphane Le Roux & Arno Pauly (2015): *Weihrauch Degrees of Finding Equilibria in Sequential Games*. In Arnold Beckmann, Victor Mitrana & Mariya Soskova, editors: *Evolving Computability, Lecture Notes in Computer Science* 9136, Springer, pp. 246–257, doi:10.1007/978-3-319-20028-6_25. Available at http://dx.doi.org/10.1007/978-3-319-20028-6_25.
- [39] Benedikt Löwe & Philipp Rohde (2002): *Games of length $\omega \cdot 2$* . *Proceedings of the AMS* 130(4), pp. 1247–1248.
- [40] Donald A. Martin (1975): *Borel Determinacy*. *Annals of Mathematics* 102(2), pp. pp. 363–371. Available at <http://www.jstor.org/stable/1971035>.
- [41] Antonio Montalbán & Richard A. Shore (2012): *The limits of determinacy in second-order arithmetic*. *Proceedings of the London Mathematical Society* 104(2), pp. 223–252, doi:10.1112/plms/pdr022. Available at <http://plms.oxfordjournals.org/content/104/2/223.abstract>.
- [42] Yiannis N. Moschovakis (1980): *Descriptive Set Theory*. *Studies in Logic and the Foundations of Mathematics* 100, North-Holland.
- [43] Uwe Mylatz (1992): *Vergleich unstetiger Funktionen in der Analysis*. Diplomarbeit, Fachbereich Informatik, FernUniversität Hagen.
- [44] Takako Nemoto (2009): *Determinacy of Wadge classes and subsystems of second order arithmetic*. *Mathematical Logic Quarterly* 55(2), pp. 154–176, doi:10.1002/malq.200710081. Available at <http://dx.doi.org/10.1002/malq.200710081>.
- [45] Takako Nemoto, MedYahya Ould MedSalem & Kazuyuki Tanaka (2007): *Infinite Games in the Cantor space and subsystems of second order arithmetic*. *Mathematical Logic Quarterly* 53(3), pp. 226–236.
- [46] Arno Pauly (2010): *How Incomputable is Finding Nash Equilibria?* *Journal of Universal Computer Science* 16(18), pp. 2686–2710, doi:10.3217/jucs-016-18-2686.
- [47] Arno Pauly (2010): *On the (semi)lattices induced by continuous reducibilities*. *Mathematical Logic Quarterly* 56(5), pp. 488–502, doi:10.1002/malq.200910104.
- [48] Arno Pauly (2012): *Computable Metamathematics and its Application to Game Theory*. Ph.D. thesis, University of Cambridge.
- [49] Arno Pauly (2012): *On the topological aspects of the theory of represented spaces*. <http://arxiv.org/abs/1204.3763>.
- [50] Arno Pauly (2015): *Computability on the countable ordinals and the Hausdorff-Kuratowski theorem*. arXiv 1501.00386.
- [51] Arno Pauly & Matthew de Brecht: *Towards Synthetic Descriptive Set Theory: An instantiation with represented spaces*. arXiv 1307.1850.
- [52] Arno Pauly & Willem Fouché (2014): *How constructive is constructing measures?* arXiv:1409.3428.
- [53] Victor L. Selivanov (2013): *Total representations*. *Logical Methods in Computer Science* 9(2).
- [54] Thorsten von Stein (1989): *Vergleich nicht konstruktiv lösbarer Probleme in der Analysis*. Diplomarbeit, Fachbereich Informatik, FernUniversität Hagen.
- [55] Klaus Weihrauch (1992): *The degrees of discontinuity of some translators between representations of the real numbers*. *Informatik Berichte* 129, FernUniversität Hagen, Hagen.
- [56] Klaus Weihrauch (1992): *The TTE-interpretation of three hierarchies of omniscience principles*. *Informatik Berichte* 130, FernUniversität Hagen, Hagen.

- [57] Martin Ziegler (2005): *Computability and continuity on the real arithmetic hierarchy and the power of type-2 nondeterminism*. In Barry S. Cooper, Benedikt Löwe & Leen Torenvliet, editors: *Proceedings of CiE 2005*, LNCS 3526, Springer, pp. 562–571.
- [58] Martin Ziegler (2007): *Real Hypercomputation and Continuity*. *Theory of Computing Systems* 41, pp. 177 – 206, doi:10.1007/s00224-006-1343-6.

Acknowledgements

This work benefited from the Royal Society International Exchange Grant IE111233 and the Marie Curie International Research Staff Exchange Scheme *Computable Analysis*, PIRSES-GA-2011- 294962.